



OPEN ACCESS

Coping with heat stress during match-play tennis: Does an individualised hydration regimen enhance performance and recovery?

Julien D Périard, Sebastien Racinais, Wade L Knez, Christopher P Herrera, Ryan J Christian, Olivier Girard

Athlete Health and Performance Research Centre, ASPETAR, Qatar Orthopaedic and Sports Medicine Hospital

Correspondence to

Dr Julien Périard, Aspetar—Qatar Orthopaedic and Sports Medicine Hospital, Research and Education Centre, PO Box 29222, Doha, Qatar; julien.periard@aspetar.com

Received 31 October 2013

Revised 17 January 2014

Accepted 19 January 2014

ABSTRACT

Objectives To determine whether an individualised hydration regimen reduces thermal, physiological and perceptual strain during match-play tennis in the heat, and minimises alterations in neuromuscular function and physical performance postmatch and into recovery.

Methods 10 men undertook two matches for an effective playing time (ball in play) of 20 min (~113 min) in ~37°C and ~33% RH conditions. Participants consumed fluids ad libitum during the first match (HOT) and followed a hydration regimen (HYD) in the second match based on undertaking play euhydrated, standardising sodium intake and minimising body mass losses.

Results HYD improved prematch urine specific gravity (1.013 ± 0.006 vs 1.021 ± 0.009 g/mL; $p < 0.05$). Body mass losses (~0.3%), fluid intake (~2 L/h) and sweat rates (~1.6 L/h) were similar between conditions. Core temperature was higher during the first 10 min of effective play in HOT ($p < 0.05$), but increased similarly (~39.3°C) on match completion. Heart rate was higher (~11 bpm) throughout HOT ($p < 0.001$). Thermal sensation was higher during the first 7.5 min of effective play in HOT ($p < 0.05$). Postmatch knee extensor and plantar flexor strength losses, along with reductions in 15 m sprint time and repeated-sprint ability ($p < 0.05$), were similar in both conditions, and were restored within 24 h.

Conclusions Both the hydration regimen and ad libitum fluid consumption allowed for minimal body mass losses (<1%). However, undertaking match-play in a euhydrated state attenuated thermal, physiological and perceptual strain. Maximal voluntary strength in the lower limbs and repeated-sprint ability deteriorated similarly in both conditions, but were restored within 24 h.

INTRODUCTION

Tennis is a multifactorial sport requiring a combination of specific physical attributes such as speed, agility, power, muscular and aerobic endurance, as well as the mental capacity to anticipate, react and make split-second decisions.¹ During prolonged (1–3 h) match-play under heat stress (>35°C), these attributes may be compromised relative to when play is undertaken in cool (15–25°C) conditions, due to the development of thermal strain (ie, a rise in core and skin temperature). Indeed, physiological (eg, heart rate) and perceptual (eg, perceived exertion and thermal comfort) responses are exacerbated during match-play in the heat, as are strength losses in the lower limbs (eg, maximal

isometric torque).^{2–3} The rise in thermal strain has also been suggested to lead to autonomic and behavioural thermoregulatory responses, whereby point duration decreases or time between points increases, in order to reduce effective playing time and overall metabolic heat load.^{3–4}

Typically, a rise in thermal strain during exercise in the heat is accompanied by progressive dehydration if fluids are not sufficiently consumed,^{5–8} which is known to exacerbate the development of hyperthermia and impair prolonged exercise performance.^{4–9–10} During match-play tennis, sweat rates vary based on ambient temperature and humidity, but can reach 2.5 L/h in certain individuals.^{11–12} It has been suggested that drinking to thirst in ecologically valid settings (eg, outdoor cycling time trial) allows athletes to compensate for these sweat rates and meet their fluid needs.^{13–14} However, the sensation of thirst may often not be perceived until 1.5 L of body water is lost,^{15–16} or a 2% body mass deficit is incurred.¹⁷ As such, ad libitum fluid consumption may lead to involuntary dehydration.¹¹ It has, therefore, been recommended that hydration regimens be individualised based on sweat rate, as well as composition (eg, sodium: Na⁺).¹ However, gastric emptying is only ~1.2 L/h,¹⁸ which may prevent some hydration regimens to fully offset fluid losses when sweating is profuse. Thus, it is further recommended to replace fluid losses to avoid dehydration >2% of pre-exercise body mass during prolonged exercise.^{5–8} It is also suggested that fluid consumption should not result in a gain of body mass; however, this latter point may be irrelevant if an individual begins exercise in a severely hypohydrated state,⁸ which is characteristic in tennis players.^{19–20} Interestingly, the regular breaks during match-play tennis allow ample opportunity to rehydrate. Despite these opportunities, it is unclear whether hydration state at the onset of and during match-play tennis in the heat influences performance and perception.

Therefore, the aim of this study was to determine whether an individualised hydration regimen reduces thermal, physiological and perceptual responses during match-play tennis in the heat, compared with the typical hydration habits of high-level players. A further aim was to determine whether implementing a hydration regimen would attenuate the level of impairment in neuromuscular function and physical performance characteristics when competing in the heat and accelerate the recovery process (24–48 h). It was hypothesised



Open Access
Scan to access more
free content



CrossMark

To cite: Périard JD, Racinais S, Knez WL, et al. *Br J Sports Med* 2014;**48**:i64–i70.

that implementing a hydration regimen based on undertaking play in a euhydrated state, minimising body mass losses and standardising Na⁺ intake during play, would prevent dehydration and attenuate the elevation in core body temperature and its concomitant deleterious influence on performance and perception.

METHODS

Subjects

Ten male tennis players (International Tennis Federation number 1–3), unacclimatised to heat, were recruited to participate in the study. The participants' mean age, height, body mass and years of match-play tennis experience were 23±5 years, 182±7.3 cm, 79±7.8 kg and 17±4 years. The participants played an average of 17±11 tournaments and 60±22 matches/year. Their weekly training volume was 12±6 h. The participants were informed of the study aims, requirements and risks before providing their written informed consent.

Study design

Participants played two simulated matches on hard-court surfaces separated by 72 h (n=4 pairs) or 144 h (n=1 pair). They were paired according to their level of play and competed against the same opponent in both matches. Both matches were played outdoors in hot conditions. In the first match (HOT: 36.8±0.3°C, 33.3±3.8% relative humidity, 34.2±0.4°C wet bulb globe temperature (WBGT)), players were instructed to eat and drink as they normally would. In the second match (HYD: 36.9±2.3°C, 32.5±12.8% relative humidity, 35.2±2.4°C WBGT), they were given a hydration regimen to follow (see Hydration section). The matches consisted of 20 min (2×10 min) of effective playing time. To calculate the effective playing time, each rally duration was measured with a stopwatch from the start (ie, ball leaving the hand of the serving player) to the end of the rally (ie, ball into the net or called out) and summed until the total duration reached 10 and 20 min (mid-match and postmatch, respectively). First serve faults and double faults were not counted for point duration. As such, total match duration (HOT: 116.6±8.1 min and HYD: 111.0±23.2 min) was of typical length for a three-set match. Data were analysed as a function of effective playing time to ensure that outcome measures were compared after an equivalent effective playing time (eg, 2.5 min) in both conditions. Each 10 min of effective play was separated by ~25 min to conduct body mass, blood lactate (see below) and physical performance measurements.²¹ One to 3 days prior to the start of the study, participants visited the testing and playing venue where they were thoroughly familiarised with the testing procedures and assessments.

Experimental protocol

On arrival on match days (9:00), participants provided a urine sample for the measurement of urine specific gravity (USG; Pal-10-S, Vitech Scientific Ltd, West Sussex, UK) and inserted a telemetric temperature pill in the rectum for measuring core temperature. After being instrumented, participants performed a standardised warm-up of running at 9 km/h for 5 min on the indoor court (~22°C). This was followed by the pre-exercise baseline neuromuscular function (see Neuromuscular function section) and physical performance assessments (see Physical performance section). These assessments were also performed post-match, and 24 and 48 h after match completion. On match days, participants were then moved outdoors and performed a 10 min tennis-specific warm-up (rallies and serves). Body mass

(Seca 769, Hamburg, Germany) and perceptual measures were recorded, along with blood lactate from a finger (Lactate Pro, Arkray Global Business Inc, Kyoto, Japan). These were also measured postmatch. Prior to starting the match, an absorbent pad with protective dressing (Tegaderm + Pad, 3M Health Care, Borken, Germany) was placed at the level of the right scapula (after cleaning the skin with deionised water) to determine sweat sodium concentration (Dimension Xpand Plus, Siemens, Munich, Germany). On the days when participants did not play, they followed a standardised training programme led by a tennis coach (~60 min) on the outdoor court after completing the assessments.

Hydration

Prior to the HOT match, participants were instructed to eat and drink as they normally would during tournament play. During the match they consumed water and a commercially available sport drink (Gatorade, Chicago, Illinois, USA) ad libitum. They were also provided with bananas and granola bars (Nature Valley, General Mills, Minneapolis, Minnesota, USA). A researcher recorded the amount and type of foods and fluids consumed during play. In addition, all participants kept a food diary to record the amount of food consumed at each meal, which were prepared for them, in order to replicate their diet before the second match.

One day prior to the HYD match, participants were asked to consume a specific volume of water (6 mL/kg body/mass) with target Na⁺ content of 185 mg/L every 2.5 h (~4–5 times/day) to ensure euhydration at the start of play. During play they were provided with and required to drink a volume of fluids that matched 70% of their sweat loss from the HOT match. This target was selected to strike a balance between allowing participant to keep changes in body mass below 2%, while considering gastrointestinal comfort and tolerance, as well as gastric emptying. The hydration regimen was based on sweat rate alone. A standardised Na⁺ content of 860 mg/L was chosen to promote the optimal maintenance of plasma Na⁺ levels and plasma volume during play, while not being too high as to negatively impact on palatability.²² The level of Na⁺ was manipulated through the use of a commercially available carbohydrate-free electrolyte supplement (Nuun & Company Inc, Seattle, Washington, USA) and sport drink (Gatorade). Although a specific volume of fluids was given to consume, participants were allowed to request more water. Additional water was supplemented accordingly to maintain the target Na⁺ content. Carbohydrate intake was matched to the HOT trial by providing the same foods. Given that the regimen was based on having played one match in the heat, the HYD trial could not be randomised.

Match-play, thermal, physiological and perceptual measurements

The scoring and timing characteristics of the matches complied with the 2012 ITF Rules²³ and are explained in a companion paper.³ Total playing time, point duration, between-point duration and effective playing percentage were calculated for each match. Three new tennis balls were used for each 10 min of effective play, with the players retrieving balls between points. Core and quadriceps skin temperature (VitalSense, Mini Mitter, Respironics, Herrsching, Germany), heart rate (Polar Electro, Lake Success, New York, USA) and perceptual measures of thermal comfort²⁴ (scales 1–7), thermal sensation²⁵ (scales 1–7) and ratings of perceived exertion²⁶ (RPE; scales 6–20) were recorded at 2.5 min intervals of effective playing time. Every

effort was made to collect these measures between games so as not to disrupt the continuity of play.

Neuromuscular function

The protocol to evaluate maximal torque production (MP35 and BSL Pro V3.6.7, Biopac Systems Inc, Santa Barbara, USA) was performed twice in a counter-balanced order, once to evaluate the knee extensors (KE) and once to evaluate the plantar flexors (PF). The neuromuscular assessment consisted of a 5 s maximal voluntary isometric contraction (MVC), on which a paired stimulus (doublet, 100 Hz) was superimposed. The MVC was followed by one doublet and three single pulses. This sequence was repeated with a 30 s rest between repetitions and 5 s between stimulations. Stimulation and measurement procedures are described in Périard *et al.*² Peak twitch torque was calculated from the mean of the six potentiated twitches. Voluntary activation (VA) was calculated with the interpolated twitch technique using the potentiated doublet as VA (%) = $[1 - (\text{superimposed doublet} / \text{resting potentiated doublet})] \times 100$.²⁷

Physical performance

Sprints

Participants performed three 15 m sprints as fast as possible separated by 15 s alongside the outdoor tennis court. Three seconds prior to the beginning of each sprint, they were asked to assume the ready position and await the start signal. Sprint times were measured on crossing the finish line using two photocells connected to an electronic timer (Microgate SRL, Bolzano, Italy). The sprints were performed in a back and forth format to allow for passive recovery during the short rest period. Initial sprint time, total sprint times and the sprint decrement score were calculated.²¹

Jumps

Squat jumps starting from a static semisquatting position ($\sim 90^\circ$ of flexion) maintained for ~ 1 s and without any preliminary movement were performed, along with countermovement jumps starting from a standing position, squatting down and then extending the knees in one continuous movement. During all jump tests, the hands were kept on the hips to minimise the contribution of the upper limbs. An electronic timer was connected to an optical acquisition system (Optojump, Microgate SRL, Bolzano, Italy) for measuring flight times (t_f). From the flight times, jump height was calculated.²⁸ The better of two trials was recorded for each jump. However, in the event that a jump was clearly not maximal or a fault was committed, it did not count as one of the two official jumps to be compared and was repeated.

Statistical analysis

All statistical calculations were performed using PASW software V21.0 (SPSS, Chicago, Illinois, USA). A repeated-measures ANOVA was performed to test significance between and within treatments. Outcome variables were tested using Mauchly's procedure for sphericity. Whenever the data violated the assumption of sphericity, *p* values and adjusted degrees of freedom based on Greenhouse-Geisser correction were reported instead. Where significant effects were established, pairwise differences were identified using the Bonferroni post hoc analysis procedure adjusted for multiple comparisons. Significant differences in hydration status were identified using Student's paired *t* tests. A non-parametric Wilcoxon matched-pairs signed-rank test was used to test differences in

match characteristics between HOT and HYD conditions. The significance level was set at $p < 0.05$. All values are expressed as means \pm SD.

RESULTS

Hydration status

The hydration regimen improved prematch USG: 1.013 ± 0.006 (HYD) vs 1.021 ± 0.009 g/mL (HOT; $p < 0.05$). Body mass was higher throughout the HYD (79.5 ± 7.9 and 79.7 ± 8.2 kg) condition compared with HOT (78.8 ± 7.8 and 78.2 ± 8.3 kg) (pre-match and postmatch, respectively; $p < 0.05$), but did not differ over time. Fluid intake during the match in HYD (2.1 ± 0.4 L/h) and HOT (1.8 ± 0.5 L/h) was similar, as were changes in body mass (HYD: 0.2 ± 1.1 vs HOT: $-0.7 \pm 1.3\%$) and sweat rates (HYD: 1.5 ± 0.3 vs HOT: 1.6 ± 0.4 L/h). Sweat $[\text{Na}^+]$ was lower after the HYD (55.8 ± 17.2 mmol/L) match compared with HOT (63.6 ± 18.7 mmol/L; $p < 0.05$), which led to a lower Na^+ loss in the HYD relative to the HOT match (2.4 ± 1.0 vs 3.0 ± 1.0 g/h; $p < 0.05$).

Match characteristics

Total match duration did not differ between the HOT and HYD conditions. Match-play characteristics: number of points ($p = 0.893$), point duration ($p = 0.786$), between-point duration ($p = 0.686$) and effective playing time ($p = 0.225$) were also similar between conditions (table 1).

Thermal, physiological and perceptual responses

Core temperature increased significantly during each 10 min segment of effective play (figure 1; $p < 0.05$); however, it was lower during the first 10 min in HYD compared with HOT ($p < 0.05$). Similar core temperatures of $39.2 \pm 0.6^\circ\text{C}$ (HYD) and $39.4 \pm 0.5^\circ\text{C}$ (HOT) were reached at match-end. Skin temperature over the thigh was similarly elevated throughout both conditions, remaining unchanged over time. Heart rate was significantly lower throughout the HYD condition compared with HOT ($p < 0.001$). Blood lactate levels were similar pre-match (1.9 ± 0.6 and 2.0 ± 0.5) and postmatch (2.6 ± 1.2 and 2.6 ± 0.6 mmol/L) in the HYD and HOT conditions, respectively. Perceptual measures are presented in figure 2. Thermal comfort and RPE increased similarly over time in both conditions ($p < 0.05$). In contrast, thermal sensation was lower up to 7.5 min of effective play in the HYD compared with HOT condition ($p < 0.05$).

Table 1 Number of points, point duration, between-point duration and effective playing percentage during 20 min of effective match-play tennis in HOT and HYD conditions

Characteristic	Condition	Match
Number of points	HOT	166.2 \pm 11.7
	HYD	166.0 \pm 29.1
Point duration (s)	HOT	7.4 \pm 0.6
	HYD	7.6 \pm 1.2
Between-point duration (s)	HOT	26.7 \pm 2.7
	HYD	23.4 \pm 5.6
Effective play (%)	HOT	16.9 \pm 1.4
	HYD	18.6 \pm 1.2

No significant difference between conditions, $p > 0.05$.

Neuromuscular function

A significant decrease in maximal voluntary KE and PF torque production was observed postmatch in both conditions relative to prematch (figure 3; $p < 0.01$). This decrease was restored within 24 h in both muscle groups. VA in the KE was lower postmatch than prematch in both conditions ($p < 0.01$), recovering within 24 h. Conversely, VA in the PF was unaffected by the conditions or exercise. Peak twitch torque in the KE and PF was lower postmatch compared with prematch (table 2; $p < 0.05$), increasing back to baseline levels at 24 and 48 h.




Figure 1 Heart rate, core and skin temperature during 20 min of effective match-play tennis in HOT and HYD conditions. †Significant difference between HOT and HYD, $p < 0.05$.

Physical performance

Initial 15 m sprint time and total sprint time increased from prematch to postmatch in both conditions (table 3; $p < 0.05$). These increases in time were restored 24 and 48 h after play in both conditions. There was no difference in the sprint decrement score between or within conditions. Performance during squat and countermovement jumps was also unaffected by the conditions or exercise.

DISCUSSION

The aim of this study was to determine whether an individualised hydration regimen—based on undertaking a tennis match




Figure 2 Thermal comfort, thermal sensation and rate of perceived exertion during 20 min of effective match-play tennis in HOT and HYD conditions. †Significant difference between HOT and HYD, $p < 0.05$.




Figure 3 Knee extensor and plantar flexor torque production and voluntary activation prior to (Pre) and following (Post, 24 h and 48 h) 20 min of effective match-play tennis in HOT and HYD conditions. *Significantly different from Pre, $p<0.05$.

(~2 h) in a euhydrated state, along with minimising body mass losses and standardising Na^+ intake—reduced thermal, physiological and perceptual responses, and minimised alterations in neuromuscular function in the lower limbs as well as physical performance parameters during and following (24–48 h) play in the heat (~37°C). Our data provide evidence that (1) the hydration regimen contributed to attenuate thermal, perceptual and physiological strain, mainly during the first hour of play, while match-play characteristics evolved similarly; (2) acute postmatch strength losses in the KE and PF, along with impairments in 15 m sprint and repeated-sprint ability (ie, total sprint time and sprint decrement score), were not influenced by prematch hydration status when body mass losses remained <1% and (3) complete recovery occurred within 24 h, regardless of the hydration strategy employed.

Hydration

The main difference conferred by the hydration regimen in the current study was that it allowed for participants to undertake the HYD match in a euhydrated state, based on USG values and a slightly higher body mass, relative to the HOT condition. The USG values noted prior to the HOT match reinforce the notion that tennis players typically have poor prematch hydration habits.^{19 20} Notwithstanding, changes in body mass during play were similar between the HOT and HYD conditions, owing to a

similar fluid intake. Although participants were instructed to follow a hydration regimen in the HYD match only, they still consumed enough fluids to maintain their body mass loss <1% in the HOT match. This is likely a reflection of the frequent breaks that occur during match-play tennis, which provide sufficient opportunities to ingest fluids and avoid excessive losses in body mass. Consequently, these losses rarely exceed 2%, even during play in the heat.^{4 19 20 29–32} In certain cases, fluid consumption even surpasses that of sweat loss.³³ Accordingly, individual sweat rates ranged from 1.0 to 2.2 (HOT) and 0.9 to 1.9 L/h (HYD) in the current study, which is similar to mean values reported in the literature under warm to hot conditions.^{12 19 31}

Despite similar sweat rates, sweat $[\text{Na}^+]$ was higher during the HOT match compared with HYD. Although sweat $[\text{Na}^+]$ in the current study was higher than previously reported during indoor (17°C)³³ and outdoor (32°C)³⁴ tennis matches, it was lower than in football players undertaking a heat-response test in 44°C conditions,³⁵ and within the range typically observed under heat stress.^{5 36} Because of the higher concentration, total Na^+ losses (HOT range: 1.3–4.4g/h; HYD range: 1.1–4.2 g/h) were greater in the HOT than in the HYD match. This is an interesting finding, which may relate to several factors because sweat $[\text{Na}^+]$ can vary based on genetic predisposition, diet, sweat rate and heat acclimatisation state.⁵ It has also been shown that dehydration can increase sweat $[\text{Na}^+]$.³⁷ In the current study, undertaking the HOT match in a hypohydrated state could have similarly contributed to increased sweat $[\text{Na}^+]$, despite attaining a similar relative level of dehydration between matches.

Interestingly, excessive sweating with elevated $[\text{Na}^+]$ is associated with a greater incidence of muscle cramping in tennis and football players.^{34 38} To avoid muscle cramping, it is recommended to add 1500 mg/L of Na^+ to water or sports drinks consumed during play.³⁴ Despite this recommendation, we chose to provide far less Na^+ during play, which resulted in no incidences of cramping. Likewise, when subjects began play in a slightly hypohydrated state and incurred minimal body mass losses (ie, HOT match), no muscle cramping occurred.

Match-play responses and characteristics

Undertaking the HYD match in a euhydrated state appears to have contributed to reduce physiological strain, as evidenced by a lower heart rate throughout play, relative to the HOT match (figure 1). The initial hydration level also seems to have reduced thermal strain, as demonstrated by a lower body core temperature during the first 10 min of effective play (~60 min of total playing time) and a trend towards being reduced during the second 10 min segment of effective play, with a significantly lower temperature at after ~100 min of total playing time in HYD. These responses may be attributed to the attenuating influence of euhydration on the rate of rise in core temperature.^{4 9 10} Interestingly, the lower core temperature observed in the HYD match appears to have influenced thermal sensation, but not thermal comfort relative to the HOT match. Hence, participants felt equally hot, yet rated the playing environment as less hot or difficult to play in during HYD. Notwithstanding, the attenuation of thermal, physiological and perceptual strain reinforces the importance of undertaking play in a euhydrated state. This is supported by evidence for a carry-over effect on these measures of strain (eg, heart rate, core temperature and RPE) negatively impacting same-day repeated-bout exercise in the heat (eg, tournament play), despite adequate rehydration and core temperature returning to baseline between bouts.^{39 40}

Table 2 Knee extensor (KE) and plantar flexor (PF) peak twitch torque prior to (Pre) and following (Post, 24 h and 48 h) 20 min of effective match-play tennis in HOT and HYD conditions

Condition	Pre	Post	24 h	48 h
KE peak twitch torque (N m)				
HOT	63.4±7.7	58.0±8.8*	62.2±8.4	64.8±8.7
HYD	63.8±8.9	57.6±9.0*	60.7±8.3	62.6±8.9
PF peak twitch torque (N m)				
HOT	15.9±3.9	14.1±4.0*	14.5±3.3	15.7±3.2
HYD	16.3±3.3	13.8±2.8*	15.1±3.9	15.7±4.0

*Significantly lower than Pre, $p<0.05$.

Table 3 Physical performance characteristics prior to (Pre) and following (Post, 24 h and 48 h) 20 min of effective match-play tennis in HOT and HYD conditions

Characteristic	Condition	Pre	Post	24 h	48 h
15 m sprint time (s)	HOT	2.59±0.13	2.68±0.13*	2.60±0.16	2.58±0.15
	HYD	2.54±0.14	2.62±0.12*	2.56±0.16	2.59±0.14
Total sprint time (s)	HOT	7.93±0.41	8.15±0.39*	8.00±0.46	7.93±0.43
	HYD	7.88±0.42	8.09±0.38*	7.89±0.40	7.98±0.40
Sprint decrement (%)	HOT	2.3±2.5	1.5±2.0	2.6±2.2	2.4±1.5
	HYD	3.3±1.8	2.8±2.4	2.7±2.0	2.8±0.5
Squat jump (cm)	HOT	35.0±4.9	34.0±3.9	34.9±5.6	35.4±5.0
	HYD	35.1±5.1	35.5±4.6	34.6±5.7	35.8±5.5
Countermovement jump (cm)	HOT	37.9±6.2	37.5±4.5	37.2±6.1	37.5±5.8
	HYD	37.7±6.5	37.4±4.5	37.2±5.5	38.1±5.3

*Significantly different from Pre, $p < 0.05$.

Ultimately, minimising the rise in core temperature and heart rate during match-play tennis in the heat is in the best interest of the player, as this will reduce the risk of heat illness and the potential for a carry-over effect on same-day matches.

The elevation in heart rate may also be associated with differences in relative exercise intensity between conditions. Indeed, increases in core temperature during prolonged heat-stress exercise are aligned with changes in relative intensity through the development of cardiovascular drift.^{41–42} Based on the core temperatures and heart rates attained during play in the current study, it may be speculated that relative exercise intensity was higher in the HOT compared with HYD match. However, RPE scores and blood lactate levels were similar across conditions, and within standard range.^{43–45} Thus, it is difficult to determine whether a difference in relative intensity occurred.

Interestingly, Carling⁴⁶ suggested that the variability in physical performance during soccer matches is mediated by the inherent demands of the game. Similarly, match-play tennis is influenced by several factors such as playing style, playing surface, tactics, technical ability, score line and fitness, all of which can impact on the ability to self-regulate physical activity levels.⁴⁵ In the current study, these factors may have contributed to influence metabolic heat production and the heart rate values attained, despite the opponent and climatic conditions remaining constant between matches. Changes in match characteristics (eg, point duration and between point duration) represent a further potential influence on cardiorespiratory load; however, no differences were noted between conditions (table 1). As such, it is unlikely that match characteristics were of significant influence on the changes noted in physiological, thermal and perceptual strain between conditions in the early stages of the matches. Nonetheless, to better understand the overall load imposed during match-play tennis, future studies should not only utilise match characteristics, but also measure on-court performance such as distance covered, number of displacements and distance per displacement, as well as stroke-specific measurements (eg, speed of groundstrokes and serve).

Neuromuscular function

Our data indicate that maximal voluntary torque production was equally impaired in the KE and PF following match-play tennis in HOT and HYD conditions (figure 3). In the KE, the decrement was associated with the development of central (ie, reduction in VA) and peripheral (ie, decrease in peak twitch torque) fatigue (table 2). Conversely, the decrease in PF torque was primarily associated with a significant reduction in twitch

contractility, as VA levels remained elevated during all contractions. The discrepancy in central fatigue development between muscle groups has been observed following running and cycling and suggests that the maximal VA is more easily attained and maintained in the PF,^{47–49} likely due to muscle fibre composition.^{50–51} Regardless, the progressive development of fatigue in both muscle groups has the potential to considerably impact performance through reductions in the efficiency of on-court movements, as well as stroke proficiency.^{1–52} However, it appears that these impairments may represent acute adjustments, as lower limb strength was recovered within 24 h.

Although it may be speculated that the initial hypohydrated state (~1% lower body mass) in which participants undertook the HOT match could have negatively influenced prematch and postmatch torque production capacity, it has been shown that moderate dehydration (~2.5% body mass deficit), isolated from acute exercise-heat stress, does not impact on strength during a single MVC or enhance fatigability during multiple contractions.⁵¹ Hence, the reduction in strength observed postmatch in both conditions appears to reflect fatigue mainly at the skeletal muscle level. This observation is supported by previous findings of strength losses in the KE and PF during brief and sustained MVCs performed immediately following prolonged (ie, cycling) and intermittent (ie, tennis) exercise, irrespective of core temperature.^{53–58}

Physical performance

The ability to maintain explosiveness of movements is a key factor for success in tennis.⁵⁶ In the current study, squat and countermovement jumps were unaffected by playing in HOT or HYD conditions (table 3). These jumps reflect the explosive power of the lower limbs and indicate that in single one-off efforts, explosiveness may be maintained, despite impairments in neuromuscular function. This may stem from the positive dose/response relationship that exists between increases in muscle temperature and muscle power, which suggests that high-velocity movements (eg, brief explosive jumping) are more temperature dependent than low-velocity movements (eg, sustained cycling at a low cadence).⁵⁹ The maintenance of brief explosive movements may also result from a fatigue-related compensatory neuromuscular reorganisation of the movement strategy, especially in the countermovement jump, which has previously been reported to remain unchanged after match-play.^{55–60}

In contrast, a reduction in body mass of ~2.7% is known to negatively affect 5 and 10 m sprint times after 120 min of match-play tennis in a warm environment (~31°C).⁶¹ Our data indicate that, despite only modest dehydration (<1%), 15 m

sprint and total sprint times were reduced postmatch relative to prematch in both conditions, whereas the sprint decrement score was maintained (albeit slightly reduced). The similarity in sprint decrement score and the reduction in initial/total sprinting times are indicative of a fast-developing fatigue during the repeated maximal efforts, resulting from match-play, rather than dehydration. This fatigue, in association with the reduction in sprint speed, has previously been suggested to reflect neuromechanical adjustments (ie, impaired musculoskeletal stiffness regulation via reductions in the tolerance of impact forces or stretch load).⁶² Taken together with the strength losses experienced by lower limb muscles (figure 3), the progressive decrease in running speed when maximal efforts are repeated after match-play tennis may result in reduced stroke effectiveness.⁴⁴ Given the relatively rapid recovery of these parameters (within 24 h), emphasis should be placed on strategies that attenuate the development of fatigue during play through specific sprint training protocols (eg, repeated efforts at or above match speed).

Limitations

Given the need to calculate individual sweat rates following the HOT match, it was not possible to counterbalance the HOT and HYD trials. Although there may have been a learning effect in the battery of assessments performed, participants were thoroughly familiarised with all tests. As such, a learning effect was unlikely to occur. Another potential limitation is the development of heat acclimatisation occurring from the HOT to the HYD match. Indeed, it is known that decreases in heart rate, perceived exertion and renal $[\text{Na}^+]$, along with plasma volume expansion, occur in 3–6 days of acclimatisation.⁶³ In contrast, decreases in resting core temperature and sweat $[\text{Na}^+]$, as well as increases in sweat rate have been suggested to develop from the fifth day of acclimatisation onward.^{63 64} Thus, while there may have been an acclimatisation effect on measures of heart rate and perceived exertion, our findings of differences in core temperature and sweat $[\text{Na}^+]$ appear to have been the result of the hydration regimen, more specifically undertaking the HYD match in a euhydrated state, especially considering that eight players had 72 h between matches and only two had 144 h.

CONCLUSION

This study provides the first direct comparison of match-play tennis in the heat of high-level players adopting their typical hydration habits versus implementing an individualised hydration regimen. We have demonstrated that an individualised hydration regimen reduces thermal (ie, core temperature), perceptual (ie, thermal sensation) and physiological (heart rate) strain, mainly during the first hour of play, relative to ad libitum fluid consumption. These reductions appear to have been conferred by undertaking the hydration match in a euhydrated state, optimising sodium intake and minimising body mass losses; however, the development of adaptations related to heat acclimatisation on these reductions cannot be overlooked. Conversely, the hydration regimen had no impact on lower limb strength losses, alterations in physical performance and match-play characteristics, as these were similarly affected by the development of fatigue during play. In addition, body mass losses remained <1% during play in HYD and HOT, and strength and physical performance parameters were restored within 24 h of match completion.

What are the new findings?

- ▶ Following an individualised hydration regimen prior to and during match-play tennis (~2 h) in the heat (~37°C) attenuates thermal, perceptual and physiological strain, relative to playing in the same environmental conditions, but consuming fluids ad libitum before and throughout play.
- ▶ The reductions in strain are mainly observed in the first hour of play and appear to stem from undertaking the matches in a euhydrated state, optimising sodium intake and preventing large decrements in body mass.
- ▶ Lower limb strength, physical performance and match-play characteristics are similarly influenced by the development of fatigue during play in the heat, irrespective of prematch hydration state, provided that body mass losses remain <1% during play.

How might it impact on clinical practice?

- ▶ Individual player habits regarding precompetition hydration should be monitored to ensure that they undertake play in a euhydrated state, especially in the heat, in order to avoid a potential carry-over effect negatively impacting same-day performance.
- ▶ General recommendations about precompetition hydration should be established not only for the hours preceding a match, but also for 24 h, in order to ensure proper hydration.

Acknowledgements The authors thank all the participants for their participation in this study. Also thanks to the coaches, Tim Colijn and Samuel Rota, for conducting the practises during the days between matches. Finally, the authors thank Mr Stephen Gurr for his input regarding nutrition and in particular his help in developing the hydration regimen.

Contributors JDP, OG, SR, CPH, WLK and RJC contributed to the development of the project. JDP, OG, SR and RJC collected and analysed the data. JDP, OG, SR, CPH, WLK and RJC had intellectual input in drafting the manuscript. All authors gave final approval for submitting the manuscript.

Funding This project was funded by the Aspire Zone Foundation Research Grant.

Competing interests None.

Ethics approval The study was approved by the Shafallah Medical Genetics Center Ethical Research Committee and conformed to the current Declaration of Helsinki guidelines.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

REFERENCES

- 1 Hornery DJ, Farrow D, Mujika I, *et al.* Fatigue in tennis: mechanisms of fatigue and effect on performance. *Sports Med* 2007;37:199–212.
- 2 Périard JD, Girard O, Racinais S. Neuromuscular adjustments of the knee extensors and plantar flexors following match-play tennis in the heat. *Br J Sports Med* 2014;48:i45–51.

- 3 Périard JD, Racinais S, Knez WL, *et al.* Thermal, physiological and perceptual strain mediate alterations in match-play tennis under heat stress. *Br J Sports Med* 2014;48:i32–8.
- 4 Morante SM, Brotherhood JR. Autonomic and behavioural thermoregulation in tennis. *Br J Sports Med* 2008;42:679–85.
- 5 Sawka MN, Burke LM, Eichner ER, *et al.* American College of Sports Medicine position stand: exercise and fluid replacement. *Med Sci Sports Exerc* 2007;39:377–90.
- 6 Kovacs MS. Hydration and temperature in tennis—a practical review. *J Sports Sci Med* 2006;5:1–9.
- 7 Kovacs MS. A review of fluid and hydration in competitive tennis. *Int J Sports Physiol Perform* 2008;3:413–23.
- 8 Maughan RJ, Shirreffs SM. Dehydration and rehydration in competitive sport. *Scand J Med Sci Sports* 2010;20:40–7.
- 9 Chevront SN, Carter RC, Sawka MN. Fluid balance and endurance exercise performance. *Curr Sports Med Rep* 2003;2:202–8.
- 10 Sawka MN, Latzka WA, Matott RP, *et al.* Hydration effects on temperature regulation. *Int J Sports Med* 1998;19:S108–10.
- 11 Bergeron MF, Armstrong LE, Maresh CM. Fluid and electrolyte losses during tennis in the heat. *Clin Sports Med* 1995;14:23–32.
- 12 Bergeron MF, Maresh CM, Armstrong LE, *et al.* Fluid-electrolyte balance associated with tennis match play in a hot environment. *Int J Sport Nutr* 1995;5:180–93.
- 13 Goulet EDB. Effect of exercise-induced dehydration on time-trial exercise performance: a meta-analysis. *Br J Sports Med* 2011;45:1149–56.
- 14 Sawka MN, Noakes TD. Does dehydration impair exercise performance? *Med Sci Sports Exerc* 2007;39:1209–17.
- 15 Armstrong LE, Hubbard RW, Szlyk PC, *et al.* Voluntary dehydration and electrolyte losses during prolonged exercise in the heat. *Aviat Space Environ Med* 1985;56:765–70.
- 16 Greenleaf JE. Problem: thirst, drinking behavior, and involuntary dehydration. *Med Sci Sports Exerc* 1992;24:645–56.
- 17 Adolph EF. *Physiology of man in the desert*. New York: Interscience, 1947.
- 18 Coyle EF, Montain SJ. Benefits of fluid replacement with carbohydrate during exercise. *Med Sci Sports Exerc* 1992;24:S324–30.
- 19 Bergeron MF, Waller JL, Marinik EL. Voluntary fluid intake and core temperature responses in adolescent tennis players: sports beverage versus water. *Br J Sports Med* 2006;40:406–10.
- 20 Hornery DJ, Farrow D, Mujika I, *et al.* An integrated physiological and performance profile of professional tennis. *Br J Sports Med* 2007;41:531–6.
- 21 Girard O, Christian RJ, Racinais S, *et al.* Heat stress does not exacerbate tennis-induced alterations in physical performance. *Br J Sports Med* 2014;48:i39–44.
- 22 Maughan R. Fluid and carbohydrate intake during exercise. In: Burke LM, Deakin V, eds. *Clinical sports nutrition*. North Ryde: McGraw-Hill, 2010:330–57.
- 23 International Tennis Federation. International Tennis Federation Rules of Tennis 2012. <http://www.itftennis.com>. 2012. <http://www.itftennis.com/media/107013/107013.pdf> (accessed 9 Jul 2013).
- 24 Bedford T. The warmth factor in comfort at work: a physiological study of heating and ventilation. Industrial Health Research Board 1936; Report No. 76.
- 25 ASHRAE. *Thermal comfort conditions*. New York: ASHRAE standard, 1966.
- 26 Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc* 1982;14:377–81.
- 27 Merton PA. Voluntary strength and fatigue. *J Physiol* 1954;123:553–64.
- 28 Lehance C, Croisier JL, Bury T. Validation du système Optojump en tant qu'outil d'évaluation de la force-vitesse (puissance) des membres inférieurs. *Sci Sports* 2005;20:131–5.
- 29 Tippet ML, Stofan JR, Lacambra M, *et al.* Core temperature and sweat responses in professional women's tennis players during tournament play in the heat. *J Athl Train* 2011;46:55.
- 30 Bergeron MF, McLeod KS, Coyle JF. Core body temperature during competition in the heat: national boys' 14 s junior tennis championships. *Br J Sports Med* 2007;41:779–83.
- 31 Morante SM, Brotherhood JR. Air temperature and physiological and subjective responses during competitive singles tennis. *Br J Sports Med* 2007;41:773–8.
- 32 McCarthy PR, Thorpe RD, Williams C. Body fluid loss during competitive tennis match-play. In: Lees A, Maynard I, Hughes M, *et al.*, eds. *Science and racket sports II*. London: E & FN Spon, 1998:52–5.
- 33 Lott MJ, Galloway SD. Fluid balance and sodium losses during indoor tennis match play. *Int J Sport Nutr Exerc Metab* 2011;21:492–500.
- 34 Bergeron MF. Heat cramps—fluid and electrolyte challenges during tennis in the heat. *J Sci Med Sport* 2003;6:19–27.
- 35 Racinais S, Mohr M, Buchheit M, *et al.* Individual responses to short-term heat acclimatisation as predictors of football performance in a hot, dry environment. *Br J Sports Med* 2012;46:810–15.
- 36 Sawka MN, Montain SJ. Fluid and electrolyte supplementation for exercise heat stress. *Am J Clin Nutr* 2000;72:564S–72S.
- 37 Morgan RM, Patterson MJ, Nimmo MA. Acute effects of dehydration on sweat composition in men during prolonged exercise in the heat. *Acta Physiol Scand* 2004;182:37–43.
- 38 Horswill CA, Stofan JR, Lacambra M, *et al.* Sodium balance during U.S. football training in the heat: cramp-prone vs. reference players. *Int J Sports Med* 2009;30:789–94.
- 39 Bergeron MF, Laird MD, Marinik EL, *et al.* Repeated-bout exercise in the heat in young athletes: physiological strain and perceptual responses. *J Appl Physiol* 2009;106:476–85.
- 40 Sawka MN, Knowlton RG, Critz JB. Thermal and circulatory responses to repeated bouts of prolonged running. *Med Sci Sports* 1979;11:177–80.
- 41 Rowell LB. *Human circulation: regulation during physical stress*. New York: Oxford University Press, 1986.
- 42 Arngimsson SA, Stewart DJ, Borrani F, *et al.* Relation of heart rate to percent VO₂peak during submaximal exercise in the heat. *J Appl Physiol* 2003;94:1162–8.
- 43 Smekal G, von Duvillard SP, Rihacek C, *et al.* A physiological profile of tennis match play. *Med Sci Sports Exerc* 2001;33:999–1005.
- 44 Ferrauti A, Bergeron MF, Pluim BM, *et al.* Physiological responses in tennis and running with similar oxygen uptake. *Eur J Appl Physiol* 2001;85:27–33.
- 45 Fernandez J, Mendez-Villanueva A, Pluim BM. Intensity of tennis match play. *Br J Sports Med* 2006;40:387–91.
- 46 Carling C. Interpreting physical performance in professional soccer match-play: should we be more pragmatic in our approach? *Sports Med* 2013;43:655–63.
- 47 Racinais S, Girard O. Neuromuscular failure is unlikely to explain the early exercise cessation in hot ambient conditions. *Psychophysiology* 2012;49:853–65.
- 48 Martin V, Kerheve H, Messonnier LA, *et al.* Central and peripheral contributions to neuromuscular fatigue induced by a 24-h treadmill run. *J Appl Physiol* 2010;108:1224–33.
- 49 Millet GY, Tomazin K, Vergès S, *et al.* Neuromuscular consequences of an extreme mountain ultra-marathon. *PLoS ONE* 2011;22:e17059.
- 50 Behm DG, Whittle J, Button D, *et al.* Intermuscle differences in activation. *Muscle Nerve* 2002;25:236–43.
- 51 Henneman E, Clamann HP, Gillies JD, *et al.* Rank order of motoneurons within a pool: law of combination. *J Neurophysiol* 1974;37:1338–49.
- 52 Girard O, Millet GP. Neuromuscular fatigue in racquet sports. *Neurol Clin* 2008;26:181–94.
- 53 Périard JD, Tammam AH, Thompson MW. Skeletal muscle strength and endurance are maintained during moderate dehydration. *Int J Sports Med* 2012;33:607–12.
- 54 Girard O, Racinais S, Micallef JP, *et al.* Spinal modulations accompany peripheral fatigue during prolonged tennis playing. *Scand J Med Sci Sports* 2011;21:455–64.
- 55 Girard O, Lattier G, Micallef JP, *et al.* Changes in exercise characteristics, maximal voluntary contraction, and explosive strength during prolonged tennis playing. *Br J Sports Med* 2006;40:521–6.
- 56 Girard O, Lattier G, Maffiuletti NA, *et al.* Neuromuscular fatigue during a prolonged intermittent exercise: application to tennis. *J Electromyogr Kinesiol* 2008;18:1038–46.
- 57 Périard JD, Cramer MN, Chapman PG, *et al.* Neuromuscular function following prolonged intense self-paced exercise in hot climatic conditions. *Eur J Appl Physiol* 2011;111:1561–9.
- 58 Nybo L, Nielsen B. Hyperthermia and central fatigue during prolonged exercise in humans. *J Appl Physiol* 2001;91:1055–60.
- 59 Racinais S, Oksa J. Temperature and neuromuscular function. *Scand J Med Sci Sports* 2010;20:1–18.
- 60 Ojala T, Häkkinen K. Effects of the tennis tournament on players' physical performance, hormonal responses, muscle damage and recovery. *J Sports Sci Med* 2013;12:240–8.
- 61 Magal M, Webster MJ, Sistrunk LE, *et al.* Comparison of glycerol and water hydration regimens on tennis-related performance. *Med Sci Sports Exerc* 2003;35:150–6.
- 62 Girard O, Micallef J-P, Millet GP. Changes in spring-mass model characteristics during repeated running sprints. *Eur J Appl Physiol* 2011;111:125–34.
- 63 Armstrong LE, Maresh CM. The induction and decay of heat acclimatisation in trained athletes. *Sports Med* 1991;12:302–12.
- 64 Garrett AT, Rehrrer NJ, Patterson MJ. Induction and decay of short-term heat acclimation in moderately and highly trained athletes. *Sports Med* 2011;41:757–71.



Coping with heat stress during match-play tennis: Does an individualised hydration regimen enhance performance and recovery?

Julien D Périard, Sebastien Racinais, Wade L Knez, et al.

Br J Sports Med 2014 48: i64-i70

doi: 10.1136/bjsports-2013-093242

Updated information and services can be found at:

http://bjsm.bmj.com/content/48/Suppl_1/i64.full.html

These include:

References

This article cites 54 articles, 16 of which can be accessed free at:

http://bjsm.bmj.com/content/48/Suppl_1/i64.full.html#ref-list-1

Article cited in:

http://bjsm.bmj.com/content/48/Suppl_1/i64.full.html#related-urls

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (135 articles)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>